

REPORT DOCUMENTATION

AD-A236 883

Approved
0704-0188

2

| | | | |
|--|--|--|--|
| 1a. REPORT SECURITY CLASSIFICATION Unclassified | | 1b. | |
| 2a. SECURITY CLASSIFICATION AUTHORITY | | 3. | |
| 2b. DECLASSIFICATION/DOWNGRADING SCHEDULE | | Distribution unlimited | |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S) | | 5. MONITORING ORGANIZATION REPORT NUMBER(S) | |
| 6a. NAME OF PERFORMING ORGANIZATION Advanced Technology Materials Inc. | | 6b. OFFICE SYMBOL (If applicable) | |
| 7a. NAME OF MONITORING ORGANIZATION Office of Naval Research | | 7b. ADDRESS (City, State, and ZIP Code) Department of the Navy 800 N. Quincy Street Arlington, VA 22217 | |
| 8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research | | 8b. OFFICE SYMBOL (If applicable) | |
| 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-90-C-0201 | | 10. SOURCE OF FUNDING NUMBERS | |
| 11. TITLE (Include Security Classification) Volatile Barium B-Diketonate Polyether Adducts. Synthesis, Characterization and Metalorganic Chemical Vapor Deposition. | | PROGRAM ELEMENT NO. | |
| 12. PERSONAL AUTHOR(S) Robin A. Gardiner, Duncan W. Brown, Peter S. Kirilin and Arnold L Rheingold | | PROJECT NO. | |
| 13a. TYPE OF REPORT Technical | | TASK NO. | |
| 13b. TIME COVERED FROM _____ TO _____ | | WORK UNIT ACCESSION NO. | |
| 14. DATE OF REPORT (Year, Month, Day) May 31, 1991 | | 15. PAGE COUNT 33 | |
| 16. SUPPLEMENTARY NOTATION | | | |

| | | | |
|------------------|-------|-----------|---|
| 17. COSATI CODES | | | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) |
| FIELD | GROUP | SUB-GROUP | |
| | | | |
| | | | |

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

The 2,5,8,11,14-pentaoxaheptadecane (tetraglyme) adducts of barium complexes of 1,1,1,5,5,5-hexafluoro-2,4-pentanedione (H(hfacac)), 1,1,1-trifluoro-2,4-pentanedione (H(tfacac)) and 2,2,6,6-tetramethyl-3,5-heptanedione (H(thd)) have been prepared and characterized by X-ray crystallography, ¹H and ¹³C nmr spectroscopies, melting and sublimation points, and elemental analysis. The thermal stability of the complexes is related to their solid state structure in which the coordination number around the barium atom is nine with a monocapped, twisted square prism structure. Metalorganic chemical vapor deposition experiments were performed using the 1,1,1,5,5,5-hexafluoro-2,4-pentanedionato and 1,1,1-trifluoro-2,4-pentanedionato complexes; the 2,2,6,6-tetramethyl-3,5-heptanedionato complex dissociated prior to sublimation. The as-deposited barium fluoride films were characterized by scanning electron microscopy and energy dispersive x-ray spectroscopy.

| | | | |
|--|--|--|--|
| 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS | | 21. ABSTRACT SECURITY CLASSIFICATION Unclassified | |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL Daniel P. Sharkey | | 22b. TELEPHONE (Include Area Code) 203-749-1100 | |
| 22c. OFFICE SYMBOL | | | |

91-01763



91 0 11 044

OFFICE OF NAVAL RESEARCH

Office of Naval Research

Contract N00014-90-C-0201

R&T Code htsc001

Technical Report No. 4

Volatile Barium β -Diketonate Polyether Adducts.
Synthesis, Characterization and Metalorganic Chemical Vapor Deposition

by

Robin A. Gardiner, Duncan W. Brown, Peter S. Kirlin, and Arnold L. Rheingold



Submitted for Publication

in

Chemistry of Materials

Advanced Technology Materials, Inc.
7 Commerce Drive
Danbury, CT 06810

May 31, 1991

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

This document has been approved for public release and sale: its distribution is unlimited.

**Volatile, Barium β -Diketonate Polyether Adducts.
Synthesis, Characterization and Metalorganic Chemical Vapor Deposition.**

R. Gardiner, D. W. Brown, and P. S. Kirlin

Advanced Technology Materials, Inc.

7 Commerce Drive, Danbury, CT

and

A. L. Rheingold

Department of Chemistry

Univ. of Delaware, Newark, DE

Abstract

The 2,5,8,11,14-pentaoxaheptadecane (tetraglyme) adducts of barium complexes of 1,1,1,5,5,5-hexafluoro-2,4-pentanedione (H(hfacac)), 1,1,1-trifluoro-2,4-pentanedione (H[tfacac]) and 2,2,6,6-tetramethyl-3,5-heptanedione (H(thd)) have been prepared and characterized by X-ray crystallography, ^1H and ^{13}C nmr spectroscopies, melting and sublimation points, and elemental analysis. The thermal stability of the complexes is related to their solid state structure in which the coordination number around the barium atom is nine with a monocapped, twisted square prism structure. Metalorganic chemical vapor deposition experiments were performed using the 1,1,1,5,5,5-hexafluoro-2,4-pentanedionato and 1,1,1-trifluoro-2,4-pentanedionato complexes; the 2,2,6,6-tetramethyl-3,5-heptanedionato complex dissociated prior to sublimation. The as-deposited barium fluoride films were characterized by scanning electron microscopy and energy dispersive x-ray spectroscopy.

Introduction

Since the discovery of superconductivity above 30K in 1986¹ numerous research efforts have focused on thin film preparation of these materials for microelectronic applications. Metalorganic Chemical Vapor Deposition (MOCVD) has significant advantages relative to physical vapor deposition (PVD) methods in the growth of superconducting thin films; large wafers can be uniformly coated and scale up to large volumes is easily accomplished. MOCVD

also allows for the exacting control of the film microstructure and stoichiometry which is essential for the manufacture of superconducting devices with optimal and reproducible properties. Uniform films of BiSrCaCuO have already been grown over large areas (45 cm^2) by MOCVD² but these films required a high temperature ($T > 850^\circ\text{C}$) anneal to become superconducting. Plasma enhanced-MOCVD (PE-MOCVD) of high temperature superconducting thin films has been reported by several groups^{3,4,5,6} at temperatures ($570^\circ\text{C} \leq T \leq 700^\circ\text{C}$) which are compatible with standard silicon integrated circuit process technology.

The primary obstacle to the widespread application of MOCVD to the growth of high T_c superconducting oxide thin films (YBaCuO, BiCaSrCuO, TlBaCaCuO) is the absence of an effective alkali metal, in particular barium, source reagent. Barium prefers high coordination numbers, up to 12,⁷ and the lack of steric saturation of many barium organometallic complexes leads them to be strongly associated in the solid state; barium organometallics often undergo thermal decomposition before vaporization.⁸ For example, barium β -diketonates ($\text{Ba}(\text{RCOCH}_2\text{COR}')_2$ where $\text{R}, \text{R}' = \text{CH}_3$) have low solubility and thermal decomposition during sublimation is great. The use of large, sterically bulky groups ($\text{R}, \text{R}' = (\text{CH}_3)_3\text{C}$, and $\text{R} = (\text{CH}_3)_3\text{C}$, $\text{R}' = \text{C}_3\text{F}_7$) to block open barium coordination sites reduces intermolecular interactions, increases solubility even in non polar solvents, and enhances volatility of the complexes.⁹ Even with bulky ligands the barium atom is still not coordinatively saturated. The current MOCVD precursors for barium (barium 1,1,1,2,2,3,3-heptafluoro-7,7-dimethyloctane-4,6-dionate ($\text{Ba}(\text{fod})_2$), and barium 2,2,6,6-tetramethyl-3,5-heptanedionate ($\text{Ba}(\text{thd})_2$) undergo partial decomposition at the temperatures ($> 200^\circ\text{C}$) required to sustain sufficient usable vapor pressure.^{10,11,12} Furthermore, association in the gas phase¹³ leads to fluctuations in element delivery rates.

Improved stability of barium transport has been accomplished by saturating the carrier gas stream going to the $\text{Ba}(\text{thd})_2$ source with the 2,2,6,6-tetramethyl-3,5-heptanedione vapor.¹⁴ Similarly, saturating the carrier gas with oxygen or nitrogen Lewis bases can lower the required bubbler temperature.

Tetrahydrofuran (THF) lowers the reagent bubbler temperature from 300°C to 180°C ¹⁵ and transport of a number of barium β -diketonates at temperatures as low as 150°C has been achieved by adding nitrogen Lewis bases (e. g., triethylamine) to the carrier gas.¹⁶ Excess hydrocarbons can lead to the co-deposition of carbon

and compete for surface sites thus interfering with the *in-situ* deposition of the high temperature superconductors. Consequently, a mononuclear barium source reagent with a vapor pressure of ~1 Torr at 150°C which is thermally stable at its vaporization temperature would simplify MOCVD process development and increase process reproducibility and control.

Our investigations have concentrated on the the factors affecting the volatility, stability and deposition chemistry of the barium complexes which determine their potential as MOCVD source reagents. We report here the synthesis and characterization of three barium β -diketonate:tetraglyme adducts, and preliminary MOCVD film growth results.

Experimental section

General Information. Synthesis and handling of all organometallic complexes was performed under nitrogen atmosphere using standard Schlenk techniques or in a Vacuum Atmospheres inert atmosphere glove box. Tetraglyme was distilled over Na/K alloy and stored over calcium hydride. Barium hydride, 1,1,1,5,5,5-hexafluoro-2,4-pentanedione, 1,1,1-trifluoro-2,4-pentanedione (Strem Chemicals) and 2,2,6,6-tetramethyl-3,5-heptanedione (Aldrich Chemical Co.) were used as received. ^1H nmr and ^{13}C nmr spectra were measured with an IBM WP200SY spectrometer. Chemical shifts are reported in ppm on a δ scale with residual resonances from the nmr solvents serving as internal standards for the ^1H (C_6D_6 δ = 7.15 ppm) and ^{13}C (C_6D_6 δ = 128.0 ppm) spectra. Elemental analysis were obtained from Galbriath Laboratories Inc., 2323 Sycamore Drive, Knoxville, TN 37921-1750.

Synthesis

Preparation of barium bis(1,1,1,5,5,5-hexafluoro-2,4-pentanedionate)·tetraglyme adduct ($\text{Ba}(\text{hfacac})_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$) · 1. 1,1,1,5,5,5-hexafluoro-2,4-pentanedione (14.74 g, 70 mmoles) was added to a stirring suspension of barium hydride (11.36 g, 35.0 mmoles) in tetraglyme. Gas was constantly evolved throughout the addition of the β -diketone. The resulting, yellow solution was decanted from a white crystalline solid (11.3 g, 42%). The solid was then washed

with ~ 50ml of cold pentane and dried under vacuum (0.2 Torr). The final product (10.9 g, 40%, m.p. 121-125°C) was purified by vacuum sublimation at 150°C and 0.2 Torr. Anal. Calcd (found) for $\text{BaC}_{20}\text{F}_{12}\text{H}_{24}\text{O}_9$: Ba, 17.75 (17.65); C, 31.05 (31.44); F, 29.47 (28.09); H, 3.13 (3.32). NMR ^1H (C_6D_6), δ 3.04 (m, 6H), 3.15 (m, 12H), 3.30 (m, 4H), 6.28 (s, 2H) ppm. ^{13}C , δ 58.98 ($\underline{\text{CH}}$), 70.37 ($\underline{\text{CH}_2}$), 70.73 ($\underline{\text{CH}_2}$), 71.51 ($\underline{\text{CH}_3}$), 87.66 ($\underline{\text{CF}_3}$) ppm.

Preparation of barium bis(1,1,1-trifluoro-2,4-pentanedionate)-(tetraglyme) adduct ($\text{Ba}(\text{tfacac})_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$) - 2. 1,1,1-trifluoro-2,4-pentanedione (5.0 g, 32.4 mmoles) was added dropwise via syringe to a stirring slurry of barium hydride (2.26 g, 16.1 mmoles) in tetraglyme. Gas was constantly evolved throughout the addition of the β -diketone. The reaction mixture was then stirred overnight and the resulting mixture was filtered to leave a white crystalline solid and an orange filtrate. The solid was washed with cold pentane and then dried in a nitrogen purge (4.2 g, 40%). 0.95 g of the crude material was sublimed at 150°C and 0.1 Torr to yield a white solid (0.60 g, 25%, m.p. 133-136°C). Anal. Calcd (found) for $\text{BaC}_{20}\text{F}_6\text{H}_{30}\text{O}_9$: Ba, 20.63 (20.52); C, 36.08 (36.55); F, 17.12 (16.94); H, 4.54 (4.54). NMR ^1H (C_6D_6), δ 1.88 (s, 6H) 2.93 (s, 6H), 3.16 (m, 4H), 3.36 (br. s, 12H), 5.83 (s, 2H) ppm. ^{13}C , δ 29.85 ($\underline{\text{CH}_3}$), 59.17 ($\underline{\text{CH}}$), 70.01 ($\underline{\text{CH}_2}$), 70.11 ($\underline{\text{CH}_2}$), 70.50 ($\underline{\text{CH}_2}$), 71.12 ($\underline{\text{CH}_3}$), 93.28 ($\underline{\text{CF}_3}$), 194.06 ($\underline{\text{C=O}}$) ppm.

Preparation of barium bis(2,2,6,6-tetramethyl-3,5-heptanedionate)-(tetraglyme) adduct ($\text{Ba}(\text{thd})_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$) - 3. 2,2,6,6-tetramethyl-3,5-heptanedione (2.65 g, 14.0 mmoles) was added to a stirring suspension of barium hydride (1.0 g, 70 mmoles) in ~ 20ml of tetraglyme; gas evolution starting immediately after the β -diketone was added and continued throughout the addition. The reaction mixture was stirred overnight to yield a cream colored solid and a pale yellow solution. The solid was filtered, washed with cold pentane and then dried under vacuum (0.2 Torr) yielding a white solid (1.04 g, 27%, m.p. 93-97°C). Anal. Calcd (found) for $\text{BaC}_{32}\text{H}_{60}\text{O}_9$: Ba, 19.83 (19.91); C, 52.93 (49.74); H, 8.33 (7.87); O, 18.91 (18.61). NMR ^1H (C_6D_6), δ 1.39 (s, 36H) 3.10 (br. s, 6H), 3.29 (m, 4H), 3.36 (br. s, 12H), 5.85 (s, 2H) ppm. ^{13}C , δ 29.78, 29.66 ($(\underline{\text{CH}_3})_3\text{C}$), 41.10 ($(\underline{\text{CH}_3})_3\text{C}$), 59.15 ($\underline{\text{CH}}$), 70.41 ($\underline{\text{CH}_2}$), 70.59 ($\underline{\text{CH}_2}$), 71.66 ($\underline{\text{CH}_3}$), 197.50 ($\underline{\text{C=O}}$) ppm.

Vapor Pressure Measurements. The vapor pressure of 1 and 2 were measured with the apparatus illustrated in Figure 1. The sample cylinder was loaded with

1.0 g of precursor in an inert atmosphere. The cylinder was then connected to the system which was subsequently evacuated and backfilled with nitrogen three times. The entire system was then evacuated, the oven temperature was set and the system was allowed to equilibrate with valves 1, 2 and 3 open to vacuum. Valve 3 was then closed and 4 opened and a pressure reading was taken after equilibration (1-2 hrs). After each measurement valve 4 was closed and 3 opened and a background reading to be taken. The vapor pressure was calculated as the difference between the total system and background pressures. The system was calibrated using the vapor pressure curve for tetraethyleneglycol.¹⁷

Vapor Pressure Results

The vapor pressure of 1 and 2 were measured at several temperatures from 120 to 180°C. The data were fit by a simple linear least squares analysis to equation 1a and b (correlation 0.893 for equation 1a and 0.938 for 1b) which are plotted in figures 2 a and b.

$$\ln P_1 (\text{torr}) = 21.875 - 9.0273 \times 10^3 / T \quad 1a$$

$$\ln P_2 (\text{torr}) = 27.477 - 1.1989 \times 10^4 / T \quad 1b$$

The values of $\Delta H_{\text{sub}}^\circ$ of 1.09 and 1.44 kJmol⁻¹ are found for the heats of sublimation for 1 and 2, respectively¹⁸.

MOCVD Experiments. Reduced pressure MOCVD experiments were carried out in a standard horizontal hot wall reactor (Figure 3). The precursor (0.5 g) was loaded into the reagent bubbler in an inert atmosphere and connected to the system without exposing the compound to air. The system was evacuated to 0.05 Torr and bubbler temperature was raised to 175°C. The nitrogen carrier gas flow, regulated by an MKS mass flow controller, was 10 cm³ min⁻¹ and the total system pressure was maintained at 4 Torr. The reagent was transported through heated lines to the quartz reactor tube which was maintained at 650°C. A typical growth run lasted 2 hrs and the film deposited on the inner wall of the quartz tube was characterized by scanning electron microscopy (SEM) and energy dispersive x-ray analysis (EDX).

Single Crystal X-ray Analysis. The data collection was performed on a Nicolet R3m diffractometer and the computer programs and the sources of the scattering factors which were taken from the SHELXTL program library (5.1).¹⁹ Colorless crystals of **1** were obtained by reduced pressure sublimation, those of **2** and **3** by slow evaporation of a toluene solution. The crystals were mounted on glass fibers with epoxy cement. The basic crystallographic parameters obtained are listed in Table I. The unit cell parameters were obtained from the least-squares fit of 25 reflections ($20^\circ \leq \theta \leq 25^\circ$). The structure of **1** was solved by direct methods which located the barium atom. **2** and **3** were each solved using Patterson maps which located the barium atoms. The remaining non-hydrogen atoms were located from subsequent difference Fourier syntheses. All hydrogen atoms were included as idealized isotropic contributions ($d(\text{CH}) = 0.960 \text{ \AA}$; $U = 1.2U$ for attached C). For **1** and **2** all non-hydrogens atoms were refined with anisotropic thermal parameters. For **3** all atoms were refined anisotropically except in one *t*-butyl group, which is disordered, where only the carbon atoms with the greater contribution to the occupancy (77%) were refined anisotropically.

Results and Discussion

Our search for barium source reagents with improved volatility and deposition chemistry has been based on a simple premise; mononuclear barium complexes require filled coordination spheres. Research into blocking the open coordination sites using multidentate Lewis bases to reduce intermolecular interactions and increase volatility has met with some success.²⁰

N,N,N',N'-tetramethylethylenediamine has been used to complex with Group II alkoxides to promote solubility.²¹ A similar increase in solubility of barium bis(pentamethylcyclopentadienide) over barium bis(cyclopentadienide) caused by the bulky pentamethylcyclopentadienide ligand has also been reported.²²

Investigations into coordination of β -diketonates with diamines and triamines lead to the formation of a number of isolated adducts²³ which were stable at room temperature but decomposed at higher temperatures ($>100^\circ\text{C}$) to yield the free amine and the parent β -diketonate.

Adducts of Group II bis(1,1,1,5,5,5-hexafluoro-2,4-pentanedionates) with the general formula $M(\text{CF}_3\text{COCHCOCF}_3)_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_3)$ ($M = \text{Ca, Sr, Ba}$ and $n = 1$) have been prepared by Spee. Coordination of the longer polyethers ($n = 4, 5$) dramatically enhanced the volatility of the barium complexes.²⁴ For example, $\text{Ba}(\text{CF}_3\text{COCHCOCF}_3)_2$ ($\text{Ba}(\text{hfacac})_2$) is not volatile below 205°C at 10^{-5} Torr,²⁵ whereas the $\text{Ba}(\text{CF}_3\text{COCHCOCF}_3)_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$ ($\text{Ba}(\text{tfacac})_2 \cdot (\text{tetraglyme})$) adduct sublimes readily at 120°C at 10^{-2} Torr. Similarly, $\text{Ba}(\text{CH}_3\text{COCHCOCF}_3)_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$ ($\text{Ba}(\text{tfacac})_2 \cdot (\text{tetraglyme})$) sublimes at 150°C at 10^{-2} Torr. Vapor pressure measurements showed that the complexes were well behaved and were considerably more volatile than the parent β -diketonate complexes.

Although both $\text{Ba}(\text{tfacac})_2$ and $\text{Ba}(\text{hfacac})_2$ are not soluble in non-coordinating solvents, the respective tetraglyme adducts dissolve readily in benzene. The barium β -diketonate/tetraglyme complexes are not hygroscopic. Coordinated water has been reported to cause instabilities in the vapor pressure of the barium complexes because the rate of water dissociation competes favorably with complex sublimation.

ORTEP diagrams (Figure 4 a, b and c) show the average Ba-O bond distance in **1**, **2** and **3** differs only slightly, being 2.782(6), 2.788(4), 2.799(5) Å, respectively. The average distances for Ba to the β -diketonate oxygen atoms are 2.705(5), 2.691(3), and 2.683(4) Å, and the Ba to tetraglyme oxygen atoms are 2.837(7), 2.867(4), and 2.893(5) Å, respectively (Table II). In **1** the plane of Ba, O(1), O(2) is $19.3(2)^\circ$ from the plane of Ba, O(3) and O(4) and in **2** the plane of Ba, O(6), O(7) is $21.4(2)^\circ$ from the plane of Ba, O(8) and O(9). The same twisted propeller shape is seen in all three structures but in **3** the angle between the plane of Ba, O(3), O(4) and Ba, O(1) and O(2) is $29.6(2)^\circ$ (Table III). This is possibly due to the steric bulk of the tertiary butyl groups at the 1 and 5 positions and may also account for the different position of the terminal methyl groups in the tetraglyme molecule. The ^{13}C nmr spectra show an increasing upfield shift in the methylene signals of the coordinated tetraglyme from **1** > **2** > **3** away from the parent polyether. This shift is indicative of the strength of coordination showing it to be weakest in $\text{Ba}(\text{thd})_2 \cdot (\text{tetraglyme})$.

Tetraglyme coordination appears more affected by the size of the R substituents than the electronegativity of the β -diketonate itself. Van der Waals repulsion of the terminal methyl groups of the tetraglyme molecule with the tertiary butyl groups of the $\text{Ba}(\text{thd})_2$ weakens the coordination of the tetraglyme with the barium atom. $\text{Ba}(\text{thd})_2 \cdot (\text{tetraglyme})$ decomposed at 110°C (0.1 Torr) to give the parent β -diketonate and tetraglyme. Preliminary experiments have been carried out with the highly sterically hindered but highly electronegative barium bis(2,2-dimethyl-6,6,7,7,8,8,8-heptafluoro-3,5-octanedionate). The tetraglyme adduct was isolated as an impure oil which dissociated at 100°C and 10^{-2} Torr to yield the parent β -diketonate and tetraglyme. Further experiments are in progress to evaluate the relative contributions of steric and electronic effects arising from the structure of the β -diketonates to the strength of coordination of tetraglyme to the β -diketonate complexes of barium.

MOCVD Results

The increased thermal stability and vapor pressure of $\text{Ba}(\text{hfacac})_2 \cdot (\text{tetraglyme})$ and $\text{Ba}(\text{tfacac})_2 \cdot (\text{tetraglyme})$ enabled their use as MOCVD source reagents. Films were deposited at $0.2 \mu\text{h}^{-1}$ on the inner wall of a quartz tube. All films were dense (Figure 5 a and b). EDX analysis indicated that the films were barium fluoride contaminated with a trace of carbon. The carbon presumably arose from the fragmentation and incorporation of the organic ligands (Figure 6 a and b). $\text{Ba}(\text{hfacac})_2$ reacts to give barium fluoride under similar conditions and the as-deposited films grown using the tetraglyme adducts do not differ in composition. Tetraglyme did not effect the formation of barium fluoride films; the formation of barium fluoride films from the decomposition of fluorine containing β -diketonates at temperatures between 500 and 700°C is well documented.^{10,25,26}

Conclusion.

We have synthesized and characterized tetraglyme adducts of barium bis(1,1,1,5,5,5-hexafluoro-2,4-pentanedionate), barium bis(1,1,1-trifluoro-2,4-pentanedionate) and barium bis(2,2,6,6-tetramethyl-3,5-heptanedionate). The fluorine containing complexes $\text{Ba}(\text{hfacac})_2 \cdot (\text{tetraglyme})$ (vapor pressure of 0.23

Torr at 150°C) and $\text{Ba}(\text{tfacac})_2 \cdot (\text{tetraglyme})$ (vapor pressure of 0.06 Torr at 150°C) have been used to grow barium fluoride films. The decomposition of the non-fluorinated $\text{Ba}(\text{thd})_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_4\text{CH}_3)$ before sublimation is attributed to the steric hindrance of the t-butyl groups.

The presence of fluorine in the β -diketonate results in the complexes decomposing to barium fluoride. Barium fluoride is thermally stable below 1000°C and the MOCVD of *in situ* superconducting YBaCuO and TlBaCaCuO thin films requires a non-fluorine containing complex. Investigations are currently underway to synthesize stable complexes with the general formula $\text{Ba}(\text{RCOCHCOR}')_2 \cdot (\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_3)$ where R and R' are non fluorinated alkyl groups. These complexes will replace current Ba MOCVD reagents as they require lower temperatures for sublimation; are stable at their sublimation temperature and have higher vapor pressure enabling superior transport.

Acknowledgements

The authors wish to thank DARPA (N00014-90-C-0201) and NSF (ISI 9002399) for their support.

Supplementary Material Available: Listings of atomic coordinates and isotropic thermal parameters for 1, 2 and 3. Ordering information is given on any current masthead page.

References

- 1 Bednorz, J. G.; Muller, K.A. *Z. Phys. B.* 1986, 64, 189.
- 2 Hamaguchi, N.; Vigil, J.; Gardiner, R.; Kirlin, P. S. *Jpn. J. Appl. Phys.* 1990, 29, L596.
- 3 Zhao, J.; Marcy, H. O.; Tonge, L. M.; Wessels, B. W.; Marks, T. J.; Kannewurf, C. R. *Solid State Commun.* 1990, 74, 1091.
- 4 Chern, C. S.; Zhao, J.; Li, Y. Q.; Norris, P.; Kear, B.; Gallois, B. *Appl. Phys. Lett.* 1990, 57, 721.
- 5 Kobayashi, K.; Ichikawa, S.; Okada, G. *Jpn. J. Appl. Phys.* 1989, 28, L2165.
- 6 Bai, G. R.; Tao, W.; Wang, R.; Xie, L. M.; Zhang, X. K. *Appl. Phys. Lett.* 1989, 55, 194.
- 7 Wells, A. F., *Structural Inorganic Chemistry*, 5th ed.; Clarendon: Oxford, 1984, 574.
- 8 Evans, W. J. *Adv. Organomet. Chem.* 1985, 24, 131.
- 9 Berg, E. W.; Herrera, N. M. *Anal. Chim. Acta.* 1972, 60, 117.
- 10 Kirlin, P. S.; Binder, R.; Gardiner, R.; Brown, D. W. *Proc. SPIE: Processing of Films for High Tc Superconducting Electronics*, 1989, 1187, 115.
- 11 Malandrino, G.; Richeson, D. S.; Marks, T. J.; DeGroot, D. C.; Schindler, J. L.; Kannerwurf, C. R. *Appl. Phys. Lett.* 1989, 66, 444.
- 12 Nakamori, T.; Abe, H.; Kanamori, T.; Shibata, S. *Jpn. J. Appl. Phys.* 1988, 27, L1265.
- 13 Belcher, R.; Cranley, C. P.; Majer, J. R.; Stephen, W. I.; Uden, P. C. *Anal. Chim. Acta.* 1972, 60, 109.
- 14 Dickinson, P. H.; Geballe, T. H.; Sanjuro, A.; Hildenbrand, D.; Craig, G.; Zisk, M.; Collman, J. J. *J. Appl. Phys.* 1989, 66, 444.
- 11 Chaffee, C. D. ed. *Superconductor Week*, 1989, 3 (39), 3.
- 16 Barron, A. R.; Buriak, J. M.; Cheatham, L.; Gordon, R. *J. Electrochemical Soc.* 1990, 137(4), 225C.
- 17 CRC Handbook of chemistry and physics, 62nd edition, 1981, D-183.
- 18 The values of $\Delta H_{\text{sub}}^{\circ}$ were derived from the Clayperon equation

$$dP/dT = (\Delta H_{\text{sub}}^{\circ}) / (RT^2).$$
- 19 G. Sheldrick, Nicolet Corp., Madison, WI.
- 20 Matsuno, S.; Uchikawa, F.; Yoshizaki, K.; *Jpn. J. Appl. Phys.* 1990, 29, L947.
- 21 U.S Patent 4,555,498, Nov 26, 1985.
- 22 Burns, C. J.; Andersen, A. J. *Organometallic Chem.* 1987, 325, 31.
- 23 Brown, D. W.; Kirlin, P. S. *Final Report DARPA Contract # N00014-88-0531*, 1989.
- 24 Spee, C. I. M. A.; Meinema, H. A.; Tumer, K.; van der Zouwen, E. A.; Mackor, A. Materials Research Society meeting, Boston, 1989.

-
- 25 Purdy, A. P.; Berry, A. D.; Holm, R. T.; Fatemi, M.; Gaskill, K. K. *Inorg. Chem.* 1989, 28, 2799.
- 26 Zhao, J.; Marcy, H. O.; Tonge, L. M.; Marks, T. J.; Wessels, B. W.; *App. Phys. Lett.*, 1988, 53, 1750.

List of figure captions

Figure 1. Diagram of apparatus used to measure vapor pressure

Figure 2a. Vapor pressure plot for $\text{Ba}(\text{CF}_3\text{COCHCOCF}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$

Figure 2b. Vapor pressure plot for $\text{Ba}(\text{CF}_3\text{COCHCOCH}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$

Figure 3. Diagram of horizontal hot wall reactor used to deposit barium fluoride films.

Figure 4a. ORTEP plot of $\text{BaC}_{20}\text{H}_{24}\text{F}_{12}\text{O}_9$ (1). H atoms are omitted for clarity.

Figure 4b. ORTEP plot of $\text{BaC}_{20}\text{H}_{30}\text{F}_6\text{O}_9$ (2). H atoms are omitted for clarity.

Figure 4c. ORTEP plot of $\text{BaC}_{20}\text{H}_{60}\text{O}_9$ (3). H atoms are omitted for clarity.

Figure 5a. Scanning Electron Micrograph of film deposited using $\text{Ba}(\text{CF}_3\text{COCHCOCF}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$.

Figure 5b. Scanning Electron Micrograph of film deposited using $\text{Ba}(\text{CF}_3\text{COCHCOCH}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$.

Figure 6a. Energy Dispersive X-ray analysis of film deposited using $\text{Ba}(\text{CF}_3\text{COCHCOCF}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$.

Figure 6b. Energy Dispersive X-ray analysis of film deposited using $\text{Ba}(\text{CF}_3\text{COCHCOCH}_3)_2 \cdot (\text{CHO}(\text{CH}_2\text{CH}_2\text{O})\text{CH}_3)$.

Table I. Crystallographic data for 1, 2 and 3.

Table II. Bond distances (Å) for 1, 2 and 3.

Table III. Bond angles (degrees) for 1, 2 and 3.




Fig 1 - GORENCE et al




Fig 2a - GARDNER et al




Fig 2b Graph of $\ln P$ vs $1/T$







Fig. 3. Gattner et al.

10 for 3.31.70 (L) - 0.7 b13









5

25



RG 1


Ca - 600000 1.00



R92

1991 14:48:33

500 counts Disp=10.230 keV
Elapsed= 0.44
48 sec



100

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|

1
 2
 3
 4
 5

10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532

1




Table I - GARDINER et. al.

(a) Crystal Parameters

| | 1 | 2 | 3 |
|--|---------------------------|------------------------|---------------------|
| formula | $C_{20}H_{24}F_{12}O_9Ba$ | $C_{20}H_{20}F_6O_9Ba$ | $C_{32}H_{60}O_9Ba$ |
| space group | $\bar{P}1$ | $\bar{P}1$ | $\bar{P}1$ |
| crystal system | triclinic | triclinic | triclinic |
| a , Å | 9.6477(21) | 9.5655(26) | 10.6063(26) |
| b , Å | 11.1547(22) | 10.4002(29) | 14.452(3) |
| c , Å | 15.4645(31) | 15.230(4) | 15.463(4) |
| α , deg | 94.151(16) | 93.010(22) | 115.311(18) |
| β , deg | 90.730(17) | 91.702(22) | 97.739(20) |
| γ , deg | 114.458(15) | 109.631(21) | 105.155(18) |
| V , Å ³ | 1509.2(5) | 1423.3(6) | 1984.0(8) |
| Z | 2 | 2 | 2 |
| cryst. dims., mm | 0.38 x 0.40 x 0.42 | 0.38 x 0.38 x 0.40 | 0.45 x 0.22 x 0. |
| cryst color | colorless | colorless | colorless |
| $D(\text{calc})$, g/cm ³ | 1.702 | 1.553 | 1.215 |
| $\mu(\text{MoK}\alpha)$, cm ⁻¹ | 14.61 | 15.07 | 10.61 |
| temp, °C | 23(1) | 23(1) | 23(1) |
| T(max)/T(min) | 1.085 | 1.099 | 1.013 |

(b) Data Collection

| | |
|----------------|---------------------------------------|
| diffractometer | Nicolet R3m |
| monochromator | graphite |
| scan technique | Wyckoff |
| radiation | MoK α (λ = 0.71073 Å) |

(Continued)

Table 1. (Continued)

| | | | |
|--|--------------------------|--------------------------|--------------------------|
| 2 θ scan range, deg | 4-50 | 4-57 | 4-48 |
| data collected (h,k,l) | $\pm 12, \pm 14, \pm 19$ | $\pm 13, \pm 14, \pm 21$ | $\pm 13, \pm 17, \pm 18$ |
| scan speed, deg/min | var. 5-20 | var. 5-20 | var. 10-30 |
| rflns. collected | 5539 | 7476 | 6477 |
| indpt. rflns. | 5320 | 7216 | 6217 |
| $R(\text{merg}), \%$ | 1.95 | 2.49 | 2.64 |
| indpt. rflns. obsvd. $F_o > 5\sigma(F_o)$ | 4207 | 5860 | 4909 |
| std. rflns. | 3 std/197 rflns | 3 std/197 rflns | 3std/197 rflns |
| var. in stds., % | <1 | ~2 | <1 |

(c) Refinement

| | | | |
|----------------------------------|-------|-------|-------|
| $R(F), \%$ | 4.85 | 4.30 | 4.63 |
| $R(wF), \%$ | 5.34 | 4.64 | 4.85 |
| $\Delta/\sigma(\text{max})$ | 0.086 | 0.006 | 0.056 |
| $\Delta(\rho), e\text{\AA}^{-3}$ | 0.635 | 1.321 | 1.063 |
| N_o/N_v | 11.10 | 18.03 | 12.52 |
| GOF | 1.415 | 1.273 | 1.180 |

Table II - GARDINER et. al.

| 1 | | 2 | | 3 | |
|------------|-----------|------------|-----------|-------------|-----------|
| Ba-O(1) | 2.699(4) | Ba-O(6) | 2.689(3) | Ba-O(4) | 2.682(6) |
| Ba-O(2) | 2.718(6) | Ba-O(7) | 2.695(4) | Ba-O(3) | 2.664(4) |
| Ba-O(3) | 2.703(8) | Ba-O(8) | 2.717(3) | Ba-O(1) | 2.718(4) |
| Ba-O(4) | 2.698(4) | Ba-O(9) | 2.662(3) | Ba-O(2) | 2.668(4) |
| Ba-O(6) | 2.860(6) | Ba-O(1) | 2.863(4) | Ba-O(5) | 2.860(5) |
| Ba-O(7) | 2.834(11) | Ba-O(2) | 2.883(3) | Ba-O(6) | 2.891(5) |
| Ba-O(8) | 2.844(8) | Ba-O(3) | 2.866(4) | Ba-O(7) | 2.940(6) |
| Ba-O(9) | 2.826(6) | Ba-O(4) | 2.870(7) | Ba-O(8) | 2.911(5) |
| Ba-O(5) | 2.821(7) | Ba-O(5) | 2.851(4) | Ba-O(9) | 2.863(5) |
| O(1)-C(2) | 1.224(9) | O(6)-C(2) | 1.242(9) | O(4)-C(16) | 1.244(9) |
| C(2)-C(3) | 1.382(14) | C(2)-C(3) | 1.386(9) | C(16)-C(17) | 1.400(9) |
| C(3)-C(4) | 1.345(10) | C(3)-C(4) | 1.378(7) | C(17)-C(18) | 1.388(11) |
| C(4)-O(2) | 1.233(6) | C(4)-O(7) | 1.237(6) | C(18)-O(3) | 1.250(9) |
| C(2)-C(1) | 1.550(11) | C(2)-C(1) | 1.523(8) | C(16)-C(15) | 1.541(12) |
| C(4)-C(5) | 1.532(16) | C(4)-C(5) | 1.524(11) | C(18)-C(19) | 1.538(9) |
| O(3)-C(7) | 1.229(7) | O(8)-C(7) | 1.224(5) | O(1)-C(7) | 1.241(6) |
| C(7)-C(8) | 1.385(12) | C(7)-C(8) | 1.411(8) | C(7)-C(6) | 1.407(6) |
| C(8)-C(9) | 1.387(13) | C(8)-C(9) | 1.364(8) | C(6)-C(5) | 1.403(8) |
| C(9)-O(4) | 1.252(7) | C(9)-O(9) | 1.258(5) | C(5)-O(2) | 1.245(6) |
| C(7)-C(6) | 1.490(15) | C(7)-C(6) | 1.515(9) | C(7)-C(8) | 1.536(8) |
| C(9)-C(10) | 1.521(11) | C(9)-C(10) | 1.518(7) | C(5)-C(4) | 1.547(10) |

TABLE III - GARAND *et al.*

| 1 | | 2 | | 3 | |
|--------------|----------|--------------|----------|--------------|----------|
| O(1)-Ba-O(2) | 63.7(1) | O(6)-Ba-O(7) | 64.3(1) | O(4)-Ba-O(3) | 64.50(1) |
| O(1)-Ba-O(3) | 144.8(2) | O(6)-Ba-O(8) | 144.0(1) | O(4)-Ba-O(1) | 144.1(1) |
| O(1)-Ba-O(4) | 149.2(2) | O(6)-Ba-O(9) | 147.9(1) | O(4)-Ba-O(2) | 142.7(2) |
| O(1)-Ba-O(6) | 106.4(1) | O(6)-Ba-O(1) | 81.1(1) | O(4)-Ba-O(5) | 116.3(1) |
| O(1)-Ba-O(7) | 93.0(2) | O(6)-Ba-O(2) | 75.4(1) | O(4)-Ba-O(6) | 83.9(2) |
| O(1)-Ba-O(8) | 74.0(2) | O(6)-Ba-O(3) | 74.2(1) | O(4)-Ba-O(7) | 72.0(2) |
| O(1)-Ba-O(9) | 74.0(2) | O(6)-Ba-O(4) | 92.5(1) | O(4)-Ba-O(8) | 73.6(2) |
| O(1)-Ba-O(5) | 85.4(2) | O(6)-Ba-O(5) | 108.6(1) | O(4)-Ba-O(9) | 76.4(2) |
| O(2)-Ba-O(3) | 151.2(1) | O(7)-Ba-O(8) | 151.2(1) | O(3)-Ba-O(1) | 153.3(1) |
| O(2)-Ba-O(4) | 89.5(1) | O(7)-Ba-O(9) | 88.7(1) | O(3)-Ba-O(2) | 90.0(1) |
| O(2)-Ba-O(6) | 69.7(2) | O(7)-Ba-O(1) | 71.6(1) | O(3)-Ba-O(5) | 70.0(2) |
| O(2)-Ba-O(7) | 111.5(2) | O(7)-Ba-O(2) | 118.7(1) | O(3)-Ba-O(6) | 93.6(2) |
| O(2)-Ba-O(8) | 135.7(2) | O(7)-Ba-O(3) | 136.9(1) | O(3)-Ba-O(7) | 129.4(2) |
| O(2)-Ba-O(9) | 115.8(2) | O(7)-Ba-O(4) | 111.6(1) | O(3)-Ba-O(8) | 127.4(2) |
| O(2)-Ba-O(5) | 73.1(2) | O(7)-Ba-O(5) | 70.9(1) | O(3)-Ba-O(9) | 81.7(1) |
| O(3)-Ba-O(4) | 63.9(2) | O(8)-Ba-O(9) | 64.6(1) | O(1)-Ba-O(2) | 63.7(1) |
| O(3)-Ba-O(6) | 91.8(2) | O(8)-Ba-O(1) | 108.0(1) | O(1)-Ba-O(5) | 97.4(1) |
| O(3)-Ba-O(7) | 72.0(2) | O(8)-Ba-O(2) | 86.7(1) | O(1)-Ba-O(6) | 98.7(2) |
| O(3)-Ba-O(8) | 71.2(2) | O(8)-Ba-O(3) | 70.4(1) | O(1)-Ba-O(7) | 76.7(2) |
| O(3)-Ba-O(9) | 84.6(2) | O(8)-Ba-O(4) | 73.0(1) | O(1)-Ba-O(8) | 68.7(2) |
| O(3)-Ba-O(5) | 106.1(2) | O(8)-Ba-O(5) | 91.4(1) | O(1)-Ba-O(9) | 93.8(1) |
| O(4)-Ba-O(6) | 75.4(2) | O(9)-Ba-O(1) | 73.2(1) | O(2)-Ba-O(5) | 76.1(1) |

(Continu

Table III. (Continued)

| 1 | | 2 | | 3 | |
|-----------------|----------|-----------------|----------|-------------------|----------|
| O(4)-Ba-O(7) | 111.8(2) | O(9)-Ba-O(2) | 105.1(1) | O(2)-Ba-O(6) | 127.6(2) |
| O(4)-Ba-O(8) | 134.7(2) | O(9)-Ba-O(3) | 134.4(1) | O(2)-Ba-O(7) | 140.4(2) |
| O(4)-Ba-O(9) | 107.8(2) | O(9)-Ba-O(4) | 113.8(1) | O(2)-Ba-O(8) | 105.7(1) |
| O(4)-Ba-O(5) | 71.7(2) | O(9)-Ba-O(5) | 76.6(1) | O(2)-Ba-O(9) | 72.1(2) |
| Ba-O(1)-C(2) | 136.1(5) | Ba-O(6)-C(2) | 137.5(3) | Ba-O(4)-C(16) | 139.0(4) |
| O(1)-C(2)-C(1) | 111.5(8) | O(6)-C(2)-C(1) | 115.9(6) | O(4)-C(16)-C(15) | 115.7(6) |
| O(1)-C(2)-C(3) | 128.0(6) | O(6)-C(2)-C(3) | 125.7(4) | O(4)-C(16)-C(17) | 125.0(7) |
| C(2)-C(3)-C(4) | 124.0(7) | C(2)-C(3)-C(4) | 124.6(5) | C(16)-C(17)-C(18) | 126.8(7) |
| C(1)-C(2)-C(3) | 120.5(7) | C(1)-C(2)-C(3) | 118.4(5) | C(15)-C(16)-C(17) | 119.3(7) |
| C(3)-C(4)-O(2) | 129.1(9) | C(3)-C(4)-O(7) | 129.9(6) | C(17)-C(18)-O(3) | 124.9(5) |
| C(3)-C(4)-C(5) | 119.2(7) | C(3)-C(4)-C(5) | 116.1(5) | C(17)-C(18)-C(19) | 119.9(7) |
| C(5)-C(4)-O(2) | 111.6(6) | C(5)-C(4)-O(7) | 114.0(4) | C(19)-C(18)-O(3) | 115.1(7) |
| C(4)-O(7)-Ba | 135.9(5) | C(4)-O(7)-Ba | 134.5(3) | C(18)-O(3)-Ba | 139.6(4) |
| Ba-O(3)-C(7) | 136.9(6) | Ba-O(8)-C(7) | 137.3(4) | Ba-O(1)-C(7) | 134.9(4) |
| O(3)-C(7)-C(6) | 115.7(8) | O(8)-C(7)-C(6) | 118.1(5) | O(1)-C(7)-C(8) | 117.3(5) |
| O(3)-C(7)-C(8) | 127.5(8) | O(8)-C(7)-C(8) | 124.0(5) | O(1)-C(7)-C(6) | 124.2(5) |
| C(6)-C(7)-C(8) | 116.8(6) | C(6)-C(7)-C(8) | 117.9(4) | C(8)-C(7)-C(6) | 118.5(4) |
| C(7)-C(8)-C(9) | 123.5(6) | C(7)-C(8)-C(9) | 125.6(4) | C(7)-C(6)-C(5) | 126.8(5) |
| C(8)-C(9)-C(10) | 117.1(6) | C(8)-C(9)-C(10) | 118.7(4) | C(6)-C(5)-C(4) | 119.4(5) |
| C(8)-C(9)-O(4) | 128.2(7) | C(8)-C(9)-O(9) | 129.1(4) | C(6)-C(5)-O(2) | 124.2(5) |
| C(10)-C(9)-O(4) | 114.7(7) | C(10)-C(9)-O(9) | 112.3(5) | C(4)-C(5)-O(2) | 116.3(5) |
| C(9)-O(4)-Ba | 136.5(5) | C(9)-O(9)-Ba | 134.8(3) | C(5)-O(2)-Ba | 135.0(4) |

Supplementary Material

Table A. Atomic coordinates ($\times 10^4$) and isotropic thermal parameters ($\text{\AA}^2 \times 10^3$) for $\text{BaC}_{20}\text{H}_{24}\text{F}_{12}\text{O}_9$ (1).

Table B. Atomic coordinates ($\times 10^4$) and isotropic thermal parameters ($\text{\AA}^2 \times 10^3$) for $\text{BaC}_{20}\text{H}_{30}\text{F}_6\text{O}_9$ (2).

Table C. Atomic coordinates ($\times 10^4$) and isotropic thermal parameters ($\text{\AA}^2 \times 10^3$) for $\text{BaC}_{20}\text{H}_{60}\text{O}_9$ (3).

| | x | y | z | U* |
|-------|-----------|-----------|-----------|---------|
| Ba | 4517.5(4) | 5003.1(3) | 2539.6(2) | 66.0(2) |
| O(1) | 6447(5) | 7414(4) | 3252(3) | 93(2) |
| O(2) | 3212(5) | 6541(4) | 3283(3) | 94(2) |
| O(3) | 4317(6) | 2683(4) | 1764(3) | 95(2) |
| O(4) | 1820(5) | 3432(4) | 1717(3) | 95(2) |
| O(5) | 3960(8) | 6375(6) | 1212(4) | 126(3) |
| O(6) | 2610(8) | 3700(6) | 3890(4) | 140(3) |
| O(7) | 5383(11) | 3676(10) | 3792(5) | 169(6) |
| O(8) | 7440(7) | 4956(8) | 2644(6) | 158(4) |
| O(9) | 6527(7) | 6150(8) | 1263(4) | 150(4) |
| F(1) | 8789(8) | 9695(6) | 2944(4) | 201(4) |
| F(2) | 8679(8) | 9631(7) | 4260(5) | 221(5) |
| F(3) | 7840(10) | 10784(6) | 3636(6) | 222(5) |
| F(4) | 1359(9) | 7788(9) | 3128(5) | 222(6) |
| F(5) | 2727(10) | 9328(7) | 3971(6) | 230(6) |
| F(6) | 1601(10) | 7434(8) | 4390(5) | 226(6) |
| F(7) | 4448(11) | 350(6) | 2028(4) | 219(6) |
| F(8) | 4376(15) | 463(9) | 726(5) | 321(10) |
| F(9) | 2671(12) | -667(6) | 1295(7) | 275(7) |
| F(10) | -1228(7) | 555(6) | 892(5) | 196(4) |
| F(11) | -1236(6) | 2114(8) | 1668(5) | 191(4) |
| F(12) | -570(7) | 2478(8) | 467(4) | 238(5) |
| C(1) | 7970(13) | 9623(8) | 3582(5) | 134(5) |
| C(2) | 6367(8) | 8461(6) | 3464(4) | 84(3) |
| C(3) | 5076(11) | 8673(7) | 3623(4) | 107(4) |
| C(4) | 3651(9) | 7715(6) | 3546(4) | 88(3) |
| C(5) | 2309(12) | 8070(9) | 3751(6) | 129(5) |
| C(6) | 3764(15) | 491(8) | 1379(5) | 134(6) |
| C(7) | 3295(9) | 1605(6) | 1511(4) | 85(3) |
| C(8) | 1776(10) | 1317(6) | 1328(4) | 95(4) |
| C(9) | 1173(8) | 2250(6) | 1414(4) | 81(3) |
| C(10) | -470(10) | 1817(9) | 1085(6) | 118(4) |
| C(11) | 1106(11) | 3600(11) | 3879(7) | 162(6) |
| C(12) | 2860(18) | 2865(11) | 4314(7) | 203(10) |
| C(13) | 4408(20) | 3025(14) | 4373(10) | 222(12) |
| C(14) | 6747(19) | 3782(19) | 3895(10) | 251(14) |
| C(15) | 7498(16) | 3917(15) | 3143(8) | 230(11) |
| C(16) | 8148(13) | 5285(15) | 1827(9) | 240(11) |
| C(17) | 8195(18) | 6384(15) | 1513(14) | 260(12) |
| C(18) | 6512(16) | 7162(12) | 891(10) | 217(9) |
| C(19) | 5019(16) | 6859(12) | 592(7) | 186(9) |
| C(20) | 2448(12) | 6259(10) | 1029(6) | 146(7) |

* Equivalent isotropic U defined as one third of the trace of the orthogonalized U_{ij} tensor

TABLE A - GARDINER et al.

| | x | y | z | U* |
|-------|----------|----------|-----------|---------|
| Ba | 639.2(2) | 41.0(2) | 2448.6(1) | 54.4(1) |
| O(1) | 1351(4) | -1500(4) | 3789(2) | 86(1) |
| O(2) | -1361(4) | -1113(4) | 3803(2) | 84(1) |
| O(3) | -2386(4) | -38(4) | 2346(3) | 96(2) |
| O(4) | -506(6) | 1351(7) | 1132(3) | 142(3) |
| O(5) | 2339(5) | 1507(5) | 1088(3) | 127(2) |
| O(6) | -967(4) | -2491(3) | 1784(2) | 87(1) |
| O(7) | 2227(4) | -1420(3) | 1739(2) | 83(1) |
| O(8) | 456(4) | 2381(3) | 3235(2) | 82(1) |
| O(9) | 3115(4) | 1676(3) | 3306(3) | 89(1) |
| F(1) | 3209(7) | -4127(6) | 939(5) | 258(5) |
| F(2) | 4143(8) | -2909(9) | 2001(5) | 250(5) |
| F(3) | 4218(8) | -2154(8) | 846(5) | 255(5) |
| F(4) | 6035(5) | 3085(6) | 3252(4) | 181(3) |
| F(5) | 5503(5) | 2898(7) | 4563(4) | 214(4) |
| F(6) | 5883(5) | 4740(4) | 3970(4) | 179(3) |
| C(1) | -2129(9) | -4847(6) | 1439(5) | 139(4) |
| C(2) | -748(6) | -3575(4) | 1573(3) | 84(2) |
| C(3) | 623(7) | -3706(4) | 1427(3) | 88(2) |
| C(4) | 1961(6) | -2645(5) | 1520(3) | 77(2) |
| C(5) | 3337(9) | -3013(7) | 1321(5) | 114(3) |
| C(6) | 569(8) | 4645(6) | 3675(4) | 108(3) |
| C(7) | 1312(6) | 3517(4) | 3488(3) | 72(2) |
| C(8) | 2855(6) | 3830(4) | 3645(3) | 77(2) |
| C(9) | 3500(5) | 2920(4) | 3571(3) | 72(2) |
| C(10) | 5246(6) | 3409(6) | 3836(4) | 97(2) |
| C(11) | 2879(7) | -1235(7) | 4010(4) | 121(3) |
| C(12) | 509(7) | -1765(5) | 4522(3) | 92(2) |
| C(13) | -1069(8) | -2203(7) | 4229(5) | 114(3) |
| C(14) | -2889(6) | -1391(7) | 3563(5) | 122(3) |
| C(15) | -3099(6) | -218(8) | 3160(5) | 127(3) |
| C(16) | -2685(8) | 954(7) | 1889(4) | 126(4) |
| C(17) | -2036(9) | 997(9) | 1029(5) | 149(5) |
| C(18) | 311(10) | 1963(9) | 494(6) | 167(6) |
| C(19) | 1858(10) | 2382(7) | 630(5) | 136(4) |
| C(20) | 3854(8) | 1755(8) | 1157(5) | 158(4) |

* Equivalent isotropic U defined as one third of the trace of the orthogonalized U_{ij} tensor

TABLE B- GARDINER et al

| | x | y | z | U |
|-------|-----------|-----------|-----------|----------|
| Ba | 1559.3(3) | 4490.8(3) | 1932.1(3) | 53.5(2) |
| O(1) | 3101(4) | 6461(3) | 2182(3) | 64(2)* |
| O(2) | 618(4) | 6120(3) | 2710(3) | 70(2)* |
| O(3) | -700(4) | 3098(4) | 1882(4) | 81(3)* |
| O(4) | 1846(4) | 2783(4) | 2105(4) | 88(3)* |
| O(5) | -653(5) | 3991(4) | 331(3) | 93(3)* |
| O(6) | 1336(6) | 3151(5) | -145(4) | 115(4)* |
| O(7) | 3866(5) | 4056(4) | 1189(4) | 117(4)* |
| O(8) | 4291(4) | 5275(5) | 3228(4) | 112(4)* |
| O(9) | 2080(5) | 5234(4) | 4028(3) | 101(3)* |
| C(1) | -1099(12) | 6950(12) | 3567(14) | 266(14)* |
| C(1') | -867(26) | 7663(20) | 3306(19) | 67(7) |
| C(2) | 477(18) | 8791(13) | 4077(14) | 242(17)* |
| C(2') | 1078(25) | 8771(21) | 4783(18) | 66(7) |
| C(3) | 1027(16) | 7960(17) | 5016(10) | 246(18)* |
| C(3') | -214(29) | 6836(24) | 4384(21) | 82(9) |
| C(4) | 316(7) | 7659(5) | 3960(5) | 84(4)* |
| C(5) | 1149(5) | 7126(4) | 3280(4) | 56(3)* |
| C(6) | 2415(6) | 7785(4) | 3322(4) | 65(3)* |
| C(7) | 3315(5) | 7431(4) | 2779(4) | 54(3)* |
| C(8) | 4664(5) | 8291(5) | 2942(5) | 68(3)* |
| C(9) | 5183(9) | 7881(7) | 2046(7) | 142(6)* |
| C(10) | 5663(7) | 8434(8) | 3821(7) | 135(6)* |
| C(11) | 4574(8) | 9382(6) | 3175(8) | 133(7)* |
| C(12) | 3150(11) | 1951(10) | 3085(12) | 245(13)* |
| C(13) | 2272(18) | 712(14) | 1424(11) | 325(17)* |
| C(14) | 1189(11) | 451(10) | 2543(11) | 230(14)* |
| C(15) | 1903(7) | 1247(5) | 2282(7) | 97(5)* |
| C(16) | 1136(6) | 1951(5) | 2109(5) | 73(3)* |
| C(17) | -266(6) | 1645(5) | 2002(5) | 79(4)* |
| C(18) | -1100(5) | 2202(5) | 1879(4) | 66(3)* |
| C(19) | 2622(6) | 1759(6) | 1795(6) | 86(4)* |
| C(20) | -3120(9) | 661(10) | 1716(12) | 214(13)* |
| C(21) | -3450(8) | 1684(8) | 884(7) | 134(7)* |
| C(22) | -2894(9) | 2543(12) | 2668(8) | 241(11)* |
| C(23) | -1748(7) | 4337(7) | 518(6) | 106(5)* |
| C(24) | -940(9) | 3063(7) | -564(6) | 107(5)* |
| C(25) | 152(11) | 2779(9) | -794(8) | 176(7)* |
| C(26) | 2411(11) | 2840(9) | -393(8) | 169(8)* |
| C(27) | 3577(12) | 3306(14) | 276(8) | 299(15)* |
| C(28) | 5108(10) | 4599(14) | 1888(9) | 224(13)* |
| C(29) | 5248(9) | 4996(10) | 2830(10) | 151(8)* |
| C(30) | 4432(8) | 5708(9) | 4205(7) | 148(7)* |
| C(31) | 3415(10) | 5861(11) | 4582(7) | 186(9)* |
| C(32) | 1407(11) | 4584(8) | 4347(7) | 153(7)* |

* Equivalent isotropic U defined as one third of the trace of the orthogonalized U_{ij} tensor

TABLE C - GARDINER et al.